

Texas Department of State Health Services

Perinatal Hepatitis B Prevention Program

Infectious Disease Prevention Division Texas Department of State Health Services



Perinatal Hepatitis B Prevention Program (PHBPP)

- Ensure that all pregnant women are tested for hepatitis B surface antigen (HBsAg).
- Ensure reporting and tracking of HBsAg-positive pregnant women.
- Ensure that delivery facilities/hospitals receive all prenatal HBsAg lab reports prior to delivery.
- Ensure identification and management of infants born to HBsAg-positive mothers
- Ensure reporting of infants born to HBsAg-positive mothers.
- Ensure identification and case management of infants born to mothers of unknown HBsAg status.



Perinatal Hepatitis B Prevention Program (PHBPP) (continued)

- Ensure timely completion of the hepatitis B vaccine series for all infants born to HBsAg-positive mothers.
- Ensure timely completion of postvaccination serologic testing (PVST) for all infants born to HBsAg-positive women.
- Ensure vaccination of household contacts 24 months of age and younger.
- Ensure program quality, monitoring, and evaluation.

Perinatal Hepatitis B Prevention Program Staff



PHBPP Coordinator, Nurse III-Vacant

Khalilah Loggins-PS IV Khalilah.Loggins@dshs.texas.gov

Catherine Peterson-PS III
Catherine.Peterson@dshs.texas.gov

Program Email TXPeriHepB@dshs.texas.gov

Program Website:

https://www.dshs.texas.gov/immunization-unit/texas-perinatal-hepatitis-bprevention-program



PHBPP Regional Coordinators

PHB Regions 1 and 9/10

Natali Miranda <u>Natali.Miranda@dshs.texas.gov</u>

PHB Regions 2/3 and 7

Amy Wong Amy.Wong@dshs.Texas.gov

PHB Regions 4/5N, and 6/5S

Sheila Williams
Sheila.Williams.@dshs.Texas.gov

PHB Region 8

Chelsea Lopez@dshs.Texas.gov

PHB Region 11

Maria Cazares

Maria.Cazares@dshs.Texas.gov

Hepatitis B Post-Exposure Prophylaxis

First dose (and IG)

• Infants born to HBsAg-positive women should receive both HBIG and the birth dose of the hepatitis B vaccine within 12 hours of birth.

Second dose

Should be administered at least four weeks after the first dose.

Third dose

• Should be administered at least eight weeks after the second dose, 16 weeks after the first dose, and no earlier than 24 weeks of age (6 months).



Texas Department of State
Health Services

Post-Vaccination Serologic Testing

- The optimal time for post-vaccination serologic testing (PVST) in infants who completed the vaccine series on time is at nine-12 months of age.
- For all others, Providers collect PVST one to two months after the completion of the vaccine series.



Health Services

Perinatal Hepatitis B Annual Report Results*

<u>Post-Exposure Prophylaxis</u> (PEP)

- Infants with recommended PEP at birth: 512 (93.1%)
- Infants with PEP and series complete by 8 months: 376 (68.4%)
- Infants with HBIG and series complete by 12 months: 407 (74%)

No PEP

- Infants with only HBIG at birth: 1 (0.2%)
- Infants with only Hep B vaccine at birth: 20 (3.6%)
- Infants that did not get any PEP or Unknown: 17 (3.1%)

Total Outcomes

- HBIG with series complete by 12 months: 407 (74%)
- Total infants with PVST by 12/31/22: 276 (50.2%)



Texas Department of State
Health Services

Goals for the Perinatal Hepatitis B Program in 2024

- Increase the number of women enrolled in the program by continuing to educate providers on testing and reporting requirements.
- Perform timely case management activities to ensure newborn infants receive post-exposure prophylaxis, and complete the vaccine series, and PVST testing.



Health Services

Perinatal Hepatitis B Summit

When: Thursday, May 23-Friday, May 24, 2024

Where: Royal Sonesta Galleria

2222 West Loop South

Houston, TX 77027



Texas Department of State Health Services



Texas Department of State Health Services

Thank you!

January Smith, MPH
Compliance Group Manager
January.Smith@dshs.texas.gov